


# REPORT ON

## Land Use/Vegetation Cover Mapping of Wardha Valley Coalfield based on Satellite Data for the Year - 2022



Submitted to  
WESTERN COALFIELD LIMITED  
NAGPUR, MAHARASTRA



**Land Use/Vegetation Cover Mapping of Wardha Valley Coalfield  
based on Satellite Data for the Year - 2022**

**March-2023**



**Remote Sensing Cell  
Geomatics Division  
CMPDI, Ranchi**

## Document Control Sheet

(1) Job No.	RSC/564922120
(2) Publication Date	March 2023
(3) Number of Pages	38
(4) Number of Figures	8
(5) Number of Tables	11
(6) Number of Plates	2
(7) Title of Report	Vegetation cover mapping of Wardha Valley Coalfield based on satellite data for the year 2022.
(8) Aim of the Report	To prepare Land use / Vegetation cover map of Wardha Valley Coalfield on 1:50000 scale for assessing the impact of coal mining on land environment.
(9) Executing Unit	Remote Sensing Cell, Geomatics Division Central Mine Planning & Design Institute Limited, Gondwana Place, Kanke Road, Ranchi 834008
(10) User Agency	Coal India Ltd. (CIL) / Western Coalfield Ltd (WCL).
(11) Authors	Mr Tilak Mondal, Chief.Manager (Remote Sensing)
(12) Security Restriction	Restricted Circulation
(13) No. of Copies	6
(14) Distribution Statement	Official

## Contents

	<b>Page No.</b>
Document Control Sheet	ii
List of Figures	iv
List of Tables	iv
List of Plates	iv
<b>1.0 Introduction</b>	<b>1 - 4</b>
1.1 Project Reference and background	
1.2 Objectives	
1.3 Location and Accessibility	
1.4 Topography & Drainage	
1.5 Reserve Forest	
<b>2.0 Remote Sensing Concept &amp; Methodology</b>	<b>5 - 17</b>
2.1 Remote Sensing	
2.2 Electromagnetic Spectrum	
2.3 Scanning System	
2.4 Data Source	
2.5 Characteristics of Satellite/Sensor	
2.6 Data Processing	
2.6.1 Geometric Correction, rectification & geo-referencing	
2.6.2 Image enhancement	
2.6.3 Training set selection	
2.6.4 Signature generation & classification	
2.6.5 Creation / Overlay of vector database in GIS	
2.6.6 Validation of classified image	
2.6.7 Final land use / vegetation cover map preparation	
<b>3.0 Landuse / Cover Mapping</b>	<b>18- 31</b>
3.1 Introduction	
3.2 Landuse / Cover Classification	
3.3 Data Analysis & Change Detection	
3.3.1 Settlements	
3.3.2 Vegetation Cover Analysis	
3.3.3 Mining Area	
3.3.4 Agricultural Land	
3.3.5 Wasteland	
3.3.6 Water Bodies	
<b>4.0 Conclusion and Recommendations</b>	<b>32 - 33</b>
4.1 Conclusion	
4.2 Recommendations	
<b>Abbreviations &amp; Glossary</b>	<b>34 - 36</b>

## List of Figures

1.1	Location Map of Wardha Valley Coal field.	4
2.1	Remote Sensing Radiation system	5
2.2	Electromagnetic Spectrum.	6
2.3	Expanded diagram of the visible and infrared regions (upper) and microwave regions (lower) showing atmospheric windows.	7
2.4	Methodology for Land use / Cover mapping.	12
2.5	UTM-Projection Relationship.	14
2.6	Land use/Cover pattern in the year 2022	24
3.1	Year-wise Comparison of Land use/Vegetation Cover in Wardha Valley Coalfield.	26

## List of Tables

2.1	Electromagnetic spectral regions.	08
2.2	Characteristics of the satellite/sensor used in the present project work.	10
2.3	Classification Accuracy Matrix.	17
3.1	Vegetation cover / landuse classes identified in Wardha Valley Coalfield.	19
3.2	Status of land use / vegetation cover pattern in the year 2022	23
3.3	Comparative status of landuse/vegetation cover in 2019 & 2022	25
3.4	Distribution of Settlements in Wardha Valley Coalfield	27
3.5	Distribution of Vegetation Cover in Wardha Valley Coalfield	28
3.6	Distribution of Mining area in Wardha Valley Coalfield	30
3.7	Distribution of Agricultural land in Wardha Valley Coalfield	30
3.8	Distribution of Wasteland in Wardha Valley Coalfield	31

## List of Plates

List of maps/plates prepared on a scale of 1:50,000 are given below:

1. Plate No.1. HQ/REM/ 01: IRS-R2-L4X FCC of Wardha Valley Coalfield
2. Plate No.2.HQ/REM/02: IRS-R2-L4X LU/LC of Wardha Valley Coalfield

## List of Drawing

1. Land Use /Vegetation Cover map of Wardha Coalfield based on satellite data IRS R2-L4FX of the year 2022.

## Chapter 1

### Introduction

#### 1.1 Project Reference

Coal India Ltd requested CMPDI to take up the study based on remote sensing satellite data for creating the geo-environmental data base of coalfield for monitoring the impact of coal mining on land use pattern and vegetation cover. Accordingly, road map for implementation of project was submitted to Coal India Ltd for land use and vegetation cover mapping of 28 major coalfields for creating the geo-environmental data base and subsequent monitoring of impact of coal mining on land environment at regular interval of three years based on remote sensing satellite data. A work order no.CIL/WBP/Env/2009/2428 dated 29.12.2009 was issued by CIL Initially for three years. Subsequently, revised work order was issued vide letter no.CIL/WBP/ENV/2011/4706 dated 12.10.12 from coal India Ltd for the period 20-12-2013to 2016-17 which was subsequently followed by another work order vide letter no: CIL/WBP/Env/2017/DP/8477 dated 21.09.2017 from Coal India Ltd for the period 2017-18 to 2021-2022 for land reclamation monitoring of all open cast mine as well as vegetation cover mapping of 19 major coalfields including Western Coalfield as per a defined plan for monitoring the impact of mining on vegetation cover. Further, a revised work order was issued vide letter no. CIL/ENVT/2022-23/W.O/10899 dated 06.07.2022 from Coal India Limited for the period 2022-23 to 2023-24.

#### Project Background

Western Coalfield Limited (WCL) is subsidiary of coal India Ltd, dedicated for maintaining the ecological balance in the region has initiated a massive plantation programme on backfilled area, OB Dump and Waste land. The advent of high resolution, multispectral satellite data has opened a new avenue in the field of mapping and monitoring of vegetation cover. The present study has taken

up to access the impact of coal mining on land use and vegetation cover in Wardha Valley Coalfield with respect to earlier study carried out in Wardha valley coalfield in the year 2019.

## **1.2 Objectives**

The objective of present study is to prepare Land Use/Cover map of Wardha Coalfields covering an area 5225.36 Square Km on scale of 1:50000 based on satellite data of the year 2022, creating the geo-environmental data base in respect of land, vegetation cover, drainage, mining area, infrastructure etc. and regular updating of data base at a regular interval of three years to assess the impact of coal mining and other activities on land use and vegetation cover in coalfield area.

## **1.3 Location & Accessibility**

Warda Valley Coalfield covering an area of about 5225.36 sq. Km. lies in the Yavatmal and Chandrapur district of Maharashtra. It is bounded by Latitude 20° 29' 06" to 20° 48' 22" and Longitudes 79° 09' 15" to 79° 26' 39" and located in the central part of India. The coalfield area is covered under Survey of India topo-sheet no. 55L/15, 55L/16, 55P/3, 55P/4, 55P/7, 55P/8, 56I/13, 56M/1 and 56M/5 RF 1:50000. This coalfield holds a premier position in India for having the considerable share of reserve of thermal grades non-coking coal for catering the demand of coal in the western part of country. Wardha Valley coalfield is well connected by rail and road ways. Chandrapur is the central town in the coalfield which is connected with Nagpur (198 Km) in the north and Wardha (120Km) towards north-west and Kazipet (250) in the south. Chandrapur is connected also via rail with Nagpur in the north and Kazipet in the south, on the main line of South-Central Railways passing through the coalfield.


## **1.4 Topography & Drainage**

The area has almost flat to gently undulating topography developed over Precambrians, Gondwanas and Trap rocks covered with black soil and alluvium. The general slope of the area is towards south. The area is drained mainly by the Wardha, the Penganga and the Erai rivers. The north-eastern Part of the area is drained by Erai river and its tributaries whereas southern part of the area is drained by Penganga flowing along the south boundary of the coalfield

## **1.5 Reserved Forests**

The reserved forests in the Wardha Valley coalfield are Tadoba, Balharsha and Bhandak in the western side, Rajura in the southern side, Satna, Raikot, Pardi and Borgaon in the eastern side.





**Fig. 1.1: Location map of Wardha Valley Coalfield**


## Chapter 2

# Remote Sensing Concepts and Methodology

### 2.1 Remote Sensing

Remote sensing is the science and art of obtaining information about an object or area through the analysis of data acquired by a device that is not in physical contact with the object or area under investigation. The term *remote sensing* is commonly restricted to methods that employ electro-magnetic energy (such as light, heat and radio waves) as the means of detecting and measuring object characteristics.

All physical objects on the earth surface continuously emit electromagnetic radiation because of the oscillations of their atomic particles. Remote sensing is largely concerned with the measurement of electro-magnetic energy from the SUN, which is reflected, scattered or emitted by the objects on the surface of the earth. Figure 2.1 schematically illustrate the generalised processes involved in electromagnetic remote sensing of the earth resources.



## 2.2 Electromagnetic Spectrum

The electromagnetic (EM) spectrum is the continuum of energy that ranges from meters to nanometres in wavelength and travels at the speed of light. Different objects on the earth surface reflect different amounts of energy in various wavelengths of the EM spectrum.





Figure 2.2 shows the electromagnetic spectrum, which is divided on the basis of wavelength into different regions that are described in Table 2.1. The EM spectrum ranges from the very short wavelengths of the gamma-ray region to the long wavelengths of the radio region. The visible region ( $0.4\text{-}0.7 \mu\text{m}$  wavelengths) occupies only a small portion of the entire EM spectrum.

Energy reflected from the objects on the surface of the earth is recorded as a function of wavelength. During daytime, the maximum amount of energy is reflected at  $0.5 \mu\text{m}$  wavelengths, which corresponds to the green band of the visible region, and is called the *reflected energy peak* (Figure 2.2). The earth also radiates energy both day and night, with the maximum energy  $9.7 \mu\text{m}$  wavelength. This *radiant energy peak* occurs in the thermal band of the IR region (Figure 2.2).



**Figure 2.3** Expanded diagrams of the visible and infrared regions (upper) and the microwave regions (lower) showing atmospheric windows. Wavelength bands of commonly used remote sensing systems are indicated. Gases responsible for atmospheric absorption are shown.

Table 2.1 Electromagnetic spectral regions			
Region	Wavelength		Remarks
<i>Gamma ray</i>	<	0.03 nm	Incoming radiation is completely absorbed by the upper atmosphere and is not available for remote sensing.
<i>X-ray</i>	0.03 to	3.00 nm	Completely absorbed by atmosphere. Not employed in remote sensing.
<i>Ultraviolet</i>	0.03 to	0.40 $\mu\text{m}$	Incoming wavelengths less than 0.3mm are completely absorbed by Ozone in the upper atmosphere.
<i>Photographic UV band</i>	0.30 to	0.40 $\mu\text{m}$	Transmitted through atmosphere. Detectable with film and photo detectors, but atmospheric scattering is severe.
<i>Visible</i>	0.40 to	0.70 $\mu\text{m}$	Imaged with film and photo detectors. Includes reflected energy peak of earth at 0.5mm.
<i>Infrared</i>	0.70 to	100.00 $\mu\text{m}$	Interaction with matter varies with wavelength. Absorption bands separate atmospheric transmission windows.
<i>Reflected IR band</i>	0.70 to	3.00 $\mu\text{m}$	Reflected solar radiation that contains no information about thermal properties of materials. The band from 0.7-0.9mm is detectable with film and is called the <i>photographic IR band</i> .
<i>Thermal IR band</i>	3.00 to 8.00 to	5.00 $\mu\text{m}$ 14.00 $\mu\text{m}$	Principal atmospheric windows in the thermal region. Images at these wavelengths are acquired by optical-mechanical scanners and special videocon systems but not by film.
<i>Microwave</i>	0.10 to	30.00 cm	Longer wavelengths can penetrate clouds, fog and rain. Images may be acquired in the active or passive mode.
<i>Radar</i>	0.10 to	30.00 cm	Active form of microwave remote sensing. Radar images are acquired at various wavelength bands.
<i>Radio</i>	>	30.00 cm	Longest wavelength portion of electromagnetic spectrum. Some classified radars with very long wavelength operate in this region.

The earth's atmosphere absorbs energy in the gamma-ray, X-ray and most of the ultraviolet (UV) region; therefore, these regions are not used for remote sensing. Details of these regions are shown in Figure 2.3. The horizontal axes show wavelength on a logarithmic scale; the vertical axes show percent atmospheric transmission of EM energy. Wavelength regions with high transmission are called *atmospheric windows* and are used to acquire remote sensing data. Detection and measurement of the recorded energy enables identification of surface objects (by their characteristic wavelength patterns or spectral signatures), both from air-borne and space-borne platforms.

## 2.3 Scanning System

The sensing device in a remotely placed platform (aircraft/satellite) records EM radiation using a *scanning system*. In scanning system, a *sensor*, with a narrow field of view is employed; this sweeps across the terrain to produce an image. The sensor receives electromagnetic energy radiated or reflected from the terrain and converts them into signal that is recorded as numerical data. In a remote sensing satellite, multiple arrays of linear sensors are used, with each array recording simultaneously a separate band of EM energy. The array of sensors employs a spectrometer to disperse the incoming energy into a spectrum. Sensors (or *detectors*) are positioned to record specific wavelength bands of energy. The information received by the sensor is suitably manipulated and transported back to the ground receiving station. The data are reconstructed on ground into digital images. The digital image data on *magnetic/optical media* consist of picture elements arranged in regular rows and columns. The position of any picture element, *pixel*, is determined on a x-y co-ordinate system. Each pixel has a numeric value called digital number (DN) that records the intensity of electromagnetic energy measured for the ground resolution cell represented by that pixel. The range of digital numbers in an image data is controlled by the radiometric resolution of the satellite's sensor system. The digital image data are further processed to produce master images of the study area. By analysing the digital data/imagery, digitally/visually, it is possible to detect, identify and classify various objects and phenomenon on the earth surface.

Remote sensing technique (airborne/satellite) in conjunction with traditional techniques harbours in an efficient, speedy and cost-effective method for natural resource management due to its inherited capabilities of being multispectral, repetitive and synoptic areal coverage. Generation of environmental 'Data Base' on land use, soil, forest, surface and subsurface water, topography and terrain characteristics, settlement and transport network, etc., and their monitoring in near real - time is very useful for environmental management planning; this is possible only with remote sensing data.

## 2.4 Data Source

The following data are used in the present study:

- **Primary Data**

Remote Sensing Satellite data viz. Resourcesat-IRS-R2/L4FX of the year 2022 having 5.8 m. spatial resolution was used in the present study. The raw digital satellite data was obtained from NRSC, Hyderabad, on CD-ROM media.

- **Secondary Data**

- Secondary (ancillary) and ground data constitute important baseline information in remote sensing, as they improve the interpretation accuracy and reliability of remotely sensed data by enabling verification of the interpreted details and by supplementing it with the information that cannot be obtained directly from the remotely sensed data. For **Wardha Valley Coalfield**, Survey of India toposheet no. 55L/15, 55L/16, 55P/3, 55P/4, 55P/7, 55P/8, 56I/13, 56M/1 and 56M/5 as well as map showing details of location of area boundary, block boundary and road supplied by WCL were used in the study.

## 2.5 Characteristics of Satellite/Sensor

The basic properties of a satellite's sensor system can be summarised as:

- (a) Spectral coverage/resolution, i.e., band locations/width; (b) spectral dimensionality: number of bands; (c) radiometric resolution: quantisation; (d) spatial resolution/instantaneous field of view or IFOV; and (e) temporal resolution. Table 2.2 illustrates the basic properties of Resourcesat -2A satellite/sensor that was used in the present study.

**Table 2.2 Characteristics of the satellite/sensor used in the present project work**


Platform	Sensor	Spectral Bands in $\mu\text{m}$	Radiometric Resolution	Spatial Resolution	Temporal Resolution	Country
IRS-R2	L4FX	B2 0.28 - 0.31 Green	7-bit (128-grey levels)	05.8 m	24 days	India
		B3 0.25 - 0.38 Red		23.5 m		
		B4 0.27 - 0.30 NIR		23.5 m		
		B5 6.90 MIR		70.5 m		
NIR: Near Infra-Red		MIR: Middle Infra-Red				

## 2.6 Data Processing

The details of data processing carried out in the present study are shown in Figure 2.4. The processing methodology involves the following major steps:

- (a) Geometric correction, rectification and geo-referencing;
- (b) Image enhancement;
- (c) Training set selection;
- (d) Signature generation and classification;
- (e) Creation/overlay of vector database;
- (f) Validation of classified image;
- (g) Final thematic map preparation.





**Fig-2.4 –Methodology of Land Use/Vegetation Cover Analysis**

### 2.6.1 Geometric correction, rectification and geo-referencing


Inaccuracies in digital imagery may occur due to 'systematic errors' attributed to earth curvature and rotation as well as 'non-systematic errors' attributed to intermittent sensor malfunctions, etc. Systematic errors are corrected at the satellite receiving station itself while non-systematic errors/random errors are corrected in pre-processing stage.

In spite of 'System / Bulk correction' carried out at supplier end; some residual errors in respect of attitude attributes still remains even after correction. Therefore, fine tuning is required for correcting the image geometrically using ground control points (GCP).

Raw digital images contain geometric distortions, which make them unusable as maps. A map is defined as a flat representation of part of the earth's spheroidal surface that should conform to an internationally accepted type of cartographic projection, so that any measurements made on the map will be accurate with those made on the ground. Any map has two basic characteristics: (a) scale and (b) projection. While *scale* is the ratio between reduced depiction of geographical features on a map and the geographical features in the real world, *projection* is the method of transforming map information from a sphere (round Earth) to a flat (map) sheet. Therefore, it is essential to transform the digital image data from a generic co-ordinate system (i.e. from line and pixel co-ordinates) to a projected co-ordinate system. In the present study georeferencing was done with the help of Survey of India (Sol) topo-sheets so that information from various sources can be compared and integrated on a GIS platform, if required.

An understanding of the basics of projection system is required before selecting any transformation model. While maps are flat surfaces, Earth however is an irregular sphere, slightly flattened at the poles and bulging at the Equator. Map projections are systemic methods for "flattening the orange peel" in measurable ways. When transferring the Earth and its irregularities onto the plane surface of a map, the following three factors are involved: (a) geoid (b) ellipsoid and (c) projection. Figure 2.5 illustrates the relationship between these three factors. The *geoid* is the rendition of the irregular spheroidal shape of the Earth; here the variations in gravity are taken into account. The observation made on the geoid is then transferred to a regular geometric

reference surface, the *ellipsoid*. Finally, the geographical relationships of the ellipsoid (in 3-D form) are transformed into the 2-D plane of a map by a transformation process called map projection. As shown in Figure 2.5, the vast majority of projections are based upon *cones*, *cylinders* and *planes*.



In the present study, **UTM projection WGS 84 Datum** is used so as to prepare the map compatible with the Sol topo-sheets. Polyconic projection is used in Sol topo-sheets as it is best suited for small - scale mapping and larger area as well as for areas with North-South orientation (viz. India). Maps prepared using these projections are a compromise of many properties; it is neither conformal perspective nor equal area. Distances, areas and shapes are true only along central meridian. Distortion increases away from central meridian. Image transformation from generic co-ordinate system to a projected co-ordinate system was carried out using ERDAS IMAGINE v.2014 digital image processing system.

## 2.6.2 Image enhancement

To improve the interpretability of the raw data, image enhancement is necessary. Most of the digital image enhancement techniques are categorised as either point or local operations. Point operations modify the value of each pixel in the image data independently. However, local

operations modify the value of each pixel based on brightness value of neighbouring pixels. Contrast manipulations/ stretching technique based on local operation was applied on the image data using ERDAS IMAGINE 2014 version s/w. The enhanced and geocoded FCC image of Wardha Valley coalfield shown in Plate No. 1.

### **2.6.3 Training set selection**

The image data were analysed based on the interpretation keys. These keys are evolved from certain fundamental image-elements such as tone/colour, size, shape, texture, pattern, location, association and shadow. Based on the image-elements and other geo-technical elements like land form, drainage pattern and physiography; training sets were selected/identified for each land use/cover class. Field survey was carried out by taking selective traverses in order to collect the ground information (or reference data) so that training sets are selected accurately in the image. This was intended to serve as an aid for classification. Based on the variability of land use/cover condition and terrain characteristics and accessibility, 150 points were selected to generate the training sets.

### **2.6.4 Signature generation and classification**

Image classification was carried out using the maximum likelihood algorithm. The classification proceeds through the following steps: (a) calculation of statistics [i.e. signature generation] for the identified training areas, and (b) the decision boundary of maximum probability based on the mean vector, variance, covariance and correlation matrix of the pixels.

After evaluating the statistical parameters of the training sets, reliability test of training sets was conducted by measuring the statistical separation between the classes that resulted from computing divergence matrix. The overall accuracy of the classification was finally assessed with reference to ground truth data. The aerial extent of each land use class in the coalfield was determined using ERDAS IMAGINE s/w 2014 version. The classified image for the year 2022 for Wardha Valley Coalfield is shown in Plate No. 2.

### **2.6.5 Creation/overlay of vector database**

Plan showing coal block boundary are superimposed on the image as vector layer in the Arc GIS database. Road network, rail network and drainage network are also digitised on Arc GIS 10.8 version database and superimposed on the classified image.

### **2.6.6 Validation of classified image**

Ground truth survey was carried out for validation of the interpreted results from the study area. Based on the validation, classification accuracy matrix was prepared. The classification accuracy matrix is shown in Table 2.3.

Classification accuracy in case of Coal quarry, Barren OB Dump and water body was 100%. Classification accuracy in case of Dense forest, Open forest, agriculture land, waste land, scrubs between 90% to 94.45%. In case of social forestry, the classification accuracy is 91.30%. Classification accuracy for urban settlement, rural settlement and Industrial settlement varies from 80.00% to 85.71% due to poor *signature separability index* whereas classification accuracy for Sand body and Fly ash pond are 66.66% and 80.00% respectively and overall classification accuracy in case of Wardha Valley Coalfield was estimated as 90.00%.

### **2.6.7 Final land use/vegetation cover map preparation**

Final land use/vegetation cover map (Plate - 2) was printed using HP Design jet 4500 Colour Plotter. The maps are prepared on 1:50,000 scale and enclosed as drawing No. 2 along with the report. A soft copy in pdf format is also enclosed.

**Table 2.3: Classification Accuracy Matrix for Wardha Valley Coalfield**

Sl. No.	Classes in the Satellite Data	Class	Total Obsrv. Points	Land use classes as observed in the field										
				C1	C2	C3	C4	C5	C6	C7	C8	C9	C10	
1	Urban Settlement	C1	05	5										
2	Dense Forest	C2	10		8	1	1							
3	Open Forest	C3	10		1	8	1							
4	Scrubs	C4	10		1	1	7	1						
5	Social Forestry	C5	10				1	8	1					
6	Agriculture Land	C6	10					1	9					
7	Waste Upland	C7	10							10				
8	Sand Body	C8	10								10			
9	Coal Quarry	C9	10									10		
10	Water Bodies	C10	10											10
<b>Total no. of observation points</b>			<b>110</b>	05	10	10	10	10	10	10	10	10	10	10
<b>% of commission</b>				00.0	20.0	20.0	30.0	20.0	10.0	0.0	0.0	0.0	0.0	0.0
<b>% of omission</b>				00.0	20.0	20.0	30.0	20.0	10.0	0.0	0.0	0.0	0.0	0.0
<b>% of Classification Accuracy</b>				100.0	80.0	80.0	70.0	80.0	90.0	100.0	100.0	100.0	100.0	100.0
<b>Overall Accuracy (%)</b>			<b>90.000</b>											

---

## Chapter 3

### Land Use/ Vegetation Cover Mapping

#### 3.1 Introduction

Land is one of the most important natural resource on which all human activities are based. Therefore, knowledge on different type of lands as well as its spatial distribution in the form of map and statistical data is vital for its geospatial planning and management for optimal use of the land resources. In mining industry, the need for information on land use/ vegetation cover pattern has gained importance due to the all-round concern on environmental impact of mining. The information on land use/ cover inventory that includes type, spatial distribution, aerial extent, location, rate and pattern of change of each category is of paramount importance for assessing the impact of coal mining on land use/ cover.

Remote sensing data with its various spectral and spatial resolution offers comprehensive and accurate information for mapping and monitoring of land use/cover pattern, dynamics of changing pattern and trends over a period of time. By analysing the data of different cut-off dates, impact of coal mining on land use and vegetation cover can be determined.

#### 3.2 Land Use/Vegetation Cover Classification

The array of information available on land use/cover requires to be arranged or grouped under a suitable framework in order to facilitate the creation of a land use/cover database. Further, to accommodate the changing land use/cover pattern, it becomes essential to develop a standardised classification system that is not only flexible in nomenclature and definition, but also capable of incorporating information obtained from the satellite data and other different sources.

The present framework of land use/cover classification has been primarily based on the '**Manual of Nationwide Land Use/ Land Cover Mapping Using Satellite Imagery**' developed by National Remote Sensing Centre, Hyderabad. Land use map was prepared on the basis of image interpretation carried out based on the satellite data for the year 2022 for Wardha Valley coalfield and following land use/cover classes are identified (Table 3.1).

<b>Table 3.1:</b> <b>Land use/cover classes identified in Wardha Valley Coalfield</b>		
	<b>Level -I</b>	<b>Level -II</b>
1	<b>Built-Up Land</b>	1.1 Urban 1.2 Rural 1.3 Industrial
2	<b>Agricultural Land</b>	2.1 Crop Land 2.2 Fallow Land
3	<b>Forest/Vegetation Cover</b>	3.1 Dense Forest 3.2 Open Forest 3.3 Scrub 3.4 Plantation under Social Forestry 3.5 Plantation on OB Dumps 3.6 Plantation on Backfilled area
4	<b>Wasteland</b>	4.1 Waste upland with / without scrubs 4.2 Fly Ash pond 4.3 Sand Body
5	<b>Mining</b>	5.1 Coal Quarry 5.2 Area under backfilling 5.3 Water filled Quarry 5.4 Barren OB Dump 5.5 Coal Dump
6	<b>Water bodies</b>	6.1 River / Streams / Reservoir / Nala etc.


Following maps are prepared on 1:50,000 scale:

1. Plate No:1 Drawing No. HQ/REM/01 - FCC of Wardha Coalfield based on IRS-R2-L4FX data of the Year 2022, Date of pass: 03.01.2022.
2. Plate No. 2: Drawing No. HQ/REM/ 02 - Land use/Cover Map of Wardha Valley Coalfield based on IRS-R2-L4FX of the year 2022, Date of pass: 03.01.2022.




### **3.3 Data Analysis & Change Detection**

Satellite data of the year 2022 were processed using ERDAS IMAGINE 2014 image processing s/w in order to interpret the various land use/cover classes present in the study area of Wardha Valley Coalfield covering 5225.36 sq.kms. The area of each land use/cover class for Wardha Valley coalfield were calculated using ERDAS IMAGINE s/w and tabulated in Table 3.2. Comparison of various land use classes between years 2019 & 2022 are shown in the Bar Chart (Fig. 3.1).



**Plate 1 : FCC (Band 4,3,2) of Wardha Valley CF based on ESA (IRS-R2-L4FX) Data of Year – 2022**




**Plate 2 : LU / LC Map of Wardha Valley CF based on ESA (IRS-R2-L4FX) Data of Year 2022**

**TABLE – 3.2: STATUS OF LAND USE/COVER PATTERN IN WARDHA VALLEY COALFIELD DURING YEAR 2022**

LAND USE CLASSES	Year-2022	
	Area (Km <sup>2</sup> )	%
<b>SETTLEMENTS</b>		
Rural Settlements	21.91	0.42
Urban Settlements	90.71	1.74
Industrial Settlements	18.82	0.36
<b>Total Settlements</b>	<b>131.44</b>	<b>2.52</b>
<b>VEGETATION COVER</b>		
<b>FOREST</b>		
<i>Dense Forest</i>	977.31	18.70
<i>Open Forest</i>	529.13	10.13
<b>Total Forest (A)</b>	<b>1506.44</b>	<b>28.83</b>
<b>SCRUBS</b>		
<i>Scrubs (B)</i>	291.51	5.58
<b>PLANTATION</b>		
<i>Social forestry</i>	54.39	1.04
<i>Plantation on OB/Backfill</i>	41.91	0.80
<b>Total Plantation ( C )</b>	<b>96.30</b>	<b>1.84</b>
<b>Total Vegetation (A+B+C)</b>	<b>1894.25</b>	<b>36.25</b>
<b>MINING AREA</b>		
Coal Quarry	30.99	0.59
Barren OB Dump	38.27	0.73
Barren Backfilled	14.13	0.27
<b>Total Mining Area</b>	<b>83.39</b>	<b>1.60</b>
<b>AGRICULTURE</b>		
Crop Land	1199.79	22.96
Fallow Land	737.63	14.12
<b>Total Agriculture</b>	<b>1937.42</b>	<b>37.08</b>
<b>WASTELANDS</b>		
Waste land	1039.47	19.89
Sand Body	34.04	0.65
<b>Total Wasteland</b>	<b>1073.51</b>	<b>20.54</b>
<b>WATERBODIES</b>		
River, nallah, pond etc.	105.35	2.02
<b>TOTAL</b>	<b>5225.36</b>	<b>100.00</b>


Distribution of various land use classes in Wardha Valley Coalfield in the year 2022 are shown in the Pie Charts (Fig. 2.6).

Figure-2.6



**TABLE – 3.3: COMPARISON OF LAND USE/COVER PATTERN IN WARDHA VALLEY COALFIELD BETWEEN YEAR 2019 & 2022**

LAND USE CLASSES	Year-2019		Year-2022		Change w.r.t. Yr 2019		Remarks
	Area (Km <sup>2</sup> )	%	Area (Km <sup>2</sup> )	%	Area (Km <sup>2</sup> )	%	
<b>SETTLEMENTS</b>							
Rural Settlements	21.41	0.41	21.91	0.42	0.50	0.01	Increase in settlement area is due to increase in industrial activity as well as other socio-economic reasons.
Urban Settlements	89.92	1.72	90.71	1.74	0.79	0.02	
Industrial Settlements	11.79	0.23	18.82	0.36	7.03	0.13	
<b>Total Settlements</b>	<b>123.12</b>	<b>2.36</b>	<b>131.44</b>	<b>2.52</b>	8.32	0.16	
<b>VEGETATION COVER</b>							
<b>FOREST</b>							
Dense Forest	867.11	16.59	977.31	18.70	110.20	2.11	Dense forest have been increased in the year 2022. Some area of open forest have been converted into dense forest.
Open Forest	558.74	10.69	529.13	10.13	-29.61	-0.57	
<b>Total Forest (A)</b>	<b>1425.85</b>	<b>27.29</b>	<b>1506.44</b>	<b>28.83</b>	80.59	1.54	
<b>SCRUBS</b>							
Scrubs (B)	386.53	7.40	291.51	5.58	-95.02	-1.82	Some scrubby area have been converted to fallow land.
<b>PLANTATION</b>							
Social forestry	51.71	0.99	54.39	1.04	2.68	0.05	Social Forestry and plantation on OB/Backfill have been increased due to afforestation drive by Govt and WCL.
Plantation on OB/Backfill	40.74	0.78	41.91	0.80	1.17	0.02	
<b>Total Plantation ( C )</b>	<b>92.45</b>	<b>1.77</b>	<b>96.30</b>	<b>1.84</b>	3.85	0.07	
<b>Total Vegetation (A+B+C)</b>	<b>1904.83</b>	<b>36.45</b>	<b>1894.25</b>	<b>36.25</b>	-10.58	-0.20	
<b>MINING AREA</b>							
Coal Quarry	29.47	0.56	30.99	0.59	1.52	0.03	Quarry increased due to mining
Barren OB Dump	39.76	0.76	38.27	0.73	-1.49	-0.03	OB dump decreased due to plantation on it.
Barren Backfilled	7.88	0.15	14.13	0.27	6.25	0.12	Backfilled have been increased..
<b>Total Mining Area</b>	<b>77.11</b>	<b>1.48</b>	<b>83.39</b>	<b>1.60</b>	6.28	<b>0.12</b>	
<b>AGRICULTURE</b>							
Crop Land	1176.89	22.52	1199.79	22.96	22.90	0.44	Waste land and scrubby area have been converted into agricultural land.
Fallow Land	579.27	11.09	737.63	14.12	158.36	3.03	
<b>Total Agriculture</b>	<b>1756.16</b>	<b>33.61</b>	<b>1937.42</b>	<b>37.08</b>	181.26	<b>3.47</b>	
<b>WASTELANDS</b>							
Waste land	1230.35	23.55	1039.47	19.89	-190.88	-3.65	Waste land have been converted into agricultural land.
Sand Body	30.84	0.59	34.04	0.65	3.20	0.06	
<b>Total Wasteland</b>	<b>1261.19</b>	<b>24.14</b>	<b>1073.51</b>	<b>20.54</b>	-187.68	<b>-3.59</b>	
<b>WATERBODIES</b>							
River, nallah, pond etc.	102.95	1.97	105.35	2.02	2.40	0.05	
<b>TOTAL</b>	<b>5225.36</b>	<b>100.00</b>	<b>5225.36</b>	<b>100.00</b>	<b>0.00</b>	<b>0.00</b>	



**Fig. 3.1 : Year-wise Comparison of Land use / Vegetation Cover in Wardha Valley Coalfield**

### 3.3.1 Settlements

All the man-made constructions covering the land surface are included under this category. Built-up land has been further divided into rural, urban and industrial classes. In the present study, industrial settlement indicates only industrial complexes excluding residential facilities. In the year 2019 the total area covered by settlements were estimated to be 123.132 sq. km (2.36%). In year 2022 the estimated area under settlements has grown to 131.44 sq. km (2.52%). There is an increase in Settlements by 8.32 sq. km. which is about 0.16% of the total area. This increase is due to more urbanisation in mining and around small towns.

The details of the land use under this category are shown in Table 3.4 as follows:

**TABLE – 3.4**

**STATUS OF CHANGE IN SETTLEMENTS IN WARDHA VALLEY COALFIELD DURING YEAR 2019 & 2022**

LAND USE CLASSES	Year-2019		Year-2022		Change w.r.t. Yr 2019		Remarks
	Area (Km <sup>2</sup> )	%	Area (Km <sup>2</sup> )	%	Area (Km <sup>2</sup> )	%	
<b>SETTLEMENTS</b>							
Rural Settlements	21.41	0.41	21.91	0.42	0.50	0.01	Increase in settlement area is due to increase in industrial activity as well as other socio-economic reasons.
Urban Settlements	89.92	1.72	90.71	1.74	0.79	0.02	
Industrial Settlements	11.79	0.23	18.82	0.36	7.03	0.13	
<b>Total Settlements</b>	<b>123.12</b>	<b>2.36</b>	<b>131.44</b>	<b>2.52</b>	<b>8.32</b>	<b>0.16</b>	

### 3.3.2 Vegetation cover Analysis

Vegetation cover in the coalfield area comprises following five classes:

- Dense Forest
- Open Forest
- Scrubs
- Plantation on Over Burden(OB) Dumps / Backfilled area, and
- Social Forestry

There has been significant variation in the land use under the vegetation classes within the area as shown below in Table 3.5.



**TABLE – 3.5**

**STATUS OF CHANGE IN VEGETATION IN WARDHA VALLEY COALFIELD DURING YEAR 2019 & 2022**

LAND USE CLASSES	Year-2019		Year-2022		Change w.r.t. Yr 2019		Remarks
	Area (Km <sup>2</sup> )	%	Area (Km <sup>2</sup> )	%	Area (Km <sup>2</sup> )	%	
<b>VEGETATION COVER</b>							
<b>FOREST</b>							
<i>Dense Forest</i>	867.11	16.59	977.31	18.70	110.20	2.11	Dense forest have been increased in the year 2022. Some area of open forest have been converted into dense forest.
<i>Open Forest</i>	558.74	10.69	529.13	10.13	-29.61	-0.57	
<b>Total Forest (A)</b>	<b>1425.85</b>	<b>27.29</b>	<b>1506.44</b>	<b>28.83</b>	<b>80.59</b>	<b>1.54</b>	
<b>SCRUBS</b>							
<i>Scrubs (B)</i>	386.53	7.40	291.51	5.58	-95.02	-1.82	Some scrubby area have been converted to fallow land.
<b>PLANTATION</b>							
<i>Social forestry</i>	51.71	0.99	54.39	1.04	2.68	0.05	Social Forestry and plantation on OB/Backfill have been increased due to afforestation drive by Govt and WCL.
<i>Plantation on OB/Backfill</i>	40.74	0.78	41.91	0.80	1.17	0.02	
<b>Total Plantation ( C )</b>	<b>92.45</b>	<b>1.77</b>	<b>96.30</b>	<b>1.84</b>	<b>3.85</b>	<b>0.07</b>	
<b>Total Vegetation (A+B+C)</b>	<b>1904.83</b>	<b>36.45</b>	<b>1894.25</b>	<b>36.25</b>	<b>-10.58</b>	<b>-0.20</b>	

**Dense forest** – Forest having crown density of above 40% comes in this class. In the year 2019 the total area covered by dense forest were estimated to be 867.11 sq. km. (16.59%). In year 2022 the estimated area under dense forest has been 977.31 sq. km. (18.70%). There is an increase in dense forest by 110.20 sq. km as open forest being converted into dense forest which is about 2.11% of the total area.

**Open Forest** – Forest having crown density between 10% to 40% comes under this class. Open forest cover over Wardha Valley coalfield which was estimated to be 558.74 sq. km (10.69%) in 2019 has been decreased to 529.13 sq. km, i.e. 10.13 % of the coalfield area in 2022. Thus the decrease in open forest is 29.61 sq. km which is 0.57% of the total coalfield area. This reduction in open forest is attributed to conversion to dense forest.

**Scrubs** – Scrubs are vegetation with crown density less than 10%. Scrubs in the coalfield are seen to be scattered signature all over the area mixed with wastelands. There is 291.51 sq km. of scrubs, i.e. 5.58% of the coalfield area in 2022. In year 2019 the scrubs covered 386.53 sq km which were 7.40% of the coalfield area. There is a decrease of 95.02 sq. km which is 1.82% of the coalfield area. The decrease is taken place because some area of scrubs has been classified as fallow land.

**Social Forestry** – Plantation which has been carried out on wastelands, along the roadsides and colonies on green belt come under this category. Analysis of data reveals Social Forestry covers 51.71 sq. km, which is 0.99% of the coalfield area in 2019. In 2022 the area covered under social forestry was 54.39 sq. km (1.04%). There is an increase of 2.68 sq. km (0.05%). This increase is due to plantation around settlements stemmed from afforestation drive of Govt.

**Plantation over OB Dump and backfilled area** – Analysis of the data reveals that WCL has carried out significant plantation on OB dumps as well as backfilled areas during the period for maintaining the ecological balance of the area. The plantation on the OB dumps and backfilled areas were estimated to be 40.74 sq. km, i.e. 0.78% of the coalfield area in 2019. In year 2022 the plantation on OB Dumps is estimated to cover an area of 41.91 sq. km which was 0.80% of the coalfield area. There is an increase of 1.17 sq. km (0.02%) in plantation over OB dumps. This is due to plantation done on OB dumps and backfill.

### 3.3.3 Mining Area

The mining area was primarily being categorized as.

- Coal Quarry
- Barren OB Dump

To make the study more relevant and to give thrust on land reclamation, in the current study some more classes have been added as follows:

- Barren Backfilled Area
- Coal Dumps
- Water filled Quarry

In the year 2019 the coal quarry was estimated to be 29.47 sq. km (0.56%) which has increased to 30.99 sq. km (0.59%) in the year 2022. This increase is due to increase in production of coal from Open cast areas. In the year 2019 the barren OB dump was estimated to be 39.76 sq. km (0.76%) which has been decreased to 38.27 sq. km (0.73%) in the year 2022. This decrease is due to increase in plantation on OB dump. In the year 2019 the barren backfilled area was estimated to be 7.88 sq. km (0.15%) which has been increased to 14.13 sq. km (0.27%) in the year 2022. The status of land Use in the mining area over the Wardha Valley Coalfield is shown in the table 3.6 below.

**TABLE – 3.6**

**Status of change in Mining Area in Wardha Valley Coalfield during the year 2019 & 2022**

LAND USE CLASSES	Year-2019		Year-2022		Change w.r.t. Yr 2019		Remarks
	Area (Km <sup>2</sup> )	%	Area (Km <sup>2</sup> )	%	Area (Km <sup>2</sup> )	%	
<b>MINING AREA</b>							
Coal Quarry	29.47	0.56	30.99	0.59	1.52	0.03	Quarry increased due to mining
Barren OB Dump	39.76	0.76	38.27	0.73	-1.49	-0.03	OB dump decreased due to plantation on it.
Barren Backfilled	7.88	0.15	14.13	0.27	6.25	0.12	Backfilled have been increased..
<b>Total Mining Area</b>	<b>77.11</b>	<b>1.48</b>	<b>83.39</b>	<b>1.60</b>	<b>6.28</b>	<b>0.12</b>	

### 3.3.4 Agricultural Land

Land primarily used for farming and production of food, fibre and other commercial and horticultural crops falls under this category. It includes crop land (irrigated and unirrigated) and fallow land (land used for cultivation, but temporarily allowed to rest)

Crop land is 1176.89 sq. km in year 2019, which is 22.52 % of the coalfield area. In year 2022 the crop land was estimated to be 1199.79 sq. km which was 22.96% of the coalfield area. There is an increase of 22.90 sq. km which is 0.44% of the coalfield due to conversion of fallow land in crop land and waste land into agricultural land. The details are shown below in Table 3.7.

**TABLE – 3.7**

**Status of change in Agricultural land in Wardha Valley Coalfield during the year 2019 & 2022**

LAND USE CLASSES	Year-2019		Year-2022		Change w.r.t. Yr 2019		Remarks
	Area (Km <sup>2</sup> )	%	Area (Km <sup>2</sup> )	%	Area (Km <sup>2</sup> )	%	
<b>AGRICULTURE</b>							
Crop Land	1176.89	22.52	1199.79	22.96	22.90	0.44	Waste land and scrubby area have been converted into agricultural land.
Fallow Land	579.27	11.09	737.63	14.12	158.36	3.03	
<b>Total Agriculture</b>	<b>1756.16</b>	<b>33.61</b>	<b>1937.42</b>	<b>37.08</b>	<b>181.26</b>	<b>3.47</b>	

### 3.3.5 Wasteland

Wasteland is degraded and unutilised class of land which is deteriorating on account of natural causes or due to lack of appropriate water and soil management. Wasteland can result from inherent/imposed constraints such as location, environment, chemical and physical properties of the soil or financial or management constraints.

The land use pattern within the area for waste lands is shown below in Table – 3.7. Waste land was estimated to be 1230.35 sq. km (23.55%) in the year 2019. In the year of 2022, waste land is estimated to be 1039.47 sq. km (19.89%). So there is a decrease of 190.88 sq. km i.e. (3.65%) of the total coalfield area. This is due to conversion of waste land into agricultural land. The details are shown below in Table 3.8.

**TABLE – 3.8**

**Status of Change in Wastelands in Wardha Valley Coalfield during the year 2019 & 2022**

LAND USE CLASSES	Year-2019		Year-2022		Change w.r.t. Yr 2019		Remarks
	Area (Km <sup>2</sup> )	%	Area (Km <sup>2</sup> )	%	Area (Km <sup>2</sup> )	%	
<b>WASTELANDS</b>							
Waste land	1230.35	23.55	1039.47	19.89	-190.88	-3.65	Waste land have been converted into agricultural land.
Sand Body	30.84	0.59	34.04	0.65	3.20	0.06	
<b>Total Wasteland</b>	<b>1261.19</b>	<b>24.14</b>	<b>1073.51</b>	<b>20.54</b>	<b>-187.68</b>	<b>-3.59</b>	

### 3.3.6 Water bodies

It is the area of impounded water includes natural lakes, rivers/streams and manmade canal, reservoirs, tanks etc. The water bodies in the study area had been estimated to be 102.95 sq. km in year 2019, which is 1.97% of the coalfield area. In 2022 it has been estimated to be 105.35 sq. km which is 2.02% of the total area. So there is an increase of 2.40 sq. km. in water bodies which is 0.05% of the total coalfield area.

## Chapter 4

### Conclusion & Recommendation

#### 4.1 Conclusion

In the present study, land use/ vegetation cover mapping has been carried out based on Sentinel 2B satellite sensor data of March, 2019 in order to monitor the impact of coal mining on land environment which may help in formulating the mitigation measures required, if any.

Study reveals that the total area of settlements which includes urban, rural and industrial settlements in the Wardha Valley coalfields covers 131.44 km<sup>2</sup> (2.52%) area. There is an increase in settlements by 8.32 sq.km over the 2019 study primarily on account of more urbanisation in mining area and around small town. Vegetation cover which includes dense forests, open forests, scrubs, avenue plantation & plantation on over-burden dumps, covers an area of 1894.25 km<sup>2</sup> (36.25%). As compared to 2019 study there is a decrease in overall vegetation cover by 10.58 km<sup>2</sup> (0.20%) this is mainly due to decrease in scrub area. The analysis further indicates that total agricultural land which includes both crop and fallow land has increased by 181.26 km<sup>2</sup> (3.47 %) because of conversion of waste land into agricultural land due to good monsoon over the last few years in Wardha Valley coalfield. The mining area which includes coal quarry, barren OB dump, barren backfilled area, covers 83.39 km<sup>2</sup> (1.60%). As compared to 2019 there is an increase in areas under mining operations due to more production of coal. Wasteland covers 1073.51 km<sup>2</sup> (20.54%) in 2022 and 1261.19 km<sup>2</sup> (24.14%) in 2019. Waste lands have reduced because some wasteland has been converted in fallow land due to good monsoon.

The detail statistical analysis is given under Table-3.2.

## **4.2 Recommendation**

It is essential to maintain the ecological balance for sustainable development of the area together with coal mining in Wardha Valley Coalfield. It is recommended that land reclamation of the mining area should be taken up on top priority by WCL. Such study should be carried out regularly to assess the impact of coal mining on land use pattern and vegetation cover in the coalfield to formulate the remedial measures, if any, required for mitigating the adverse impact of coal mining on land environment. Such regional study will also be helpful in assessing the environmental degradation/ upgradation carried out by different industries operating in the coalfield area.

## ABBREVIATIONS

Sol	Survey of India
MoEF & CC	Ministry of Environment, Forest & Climate Change
CIL	Coal India Limited
ECL	Eastern Coalfields Limited
BCCL	Bharat Coking Coal Limited
CCL	Central Coalfields Limited
WCL	Western Coalfields Limited
SECL	South Eastern Coalfields Limited
NCL	Northern Coalfields Limited
MCL	Mahanadi Coalfields Limited
NEC	North Eastern Coalfields
CMPDIL	Central Mine Planning & Design Institute Ltd
NRSC	National Remote Sensing Centre
R2/ R2A	ResourceSat Satellites
LISS - 4	Linear Imaging and Self Scanning Sensor
FCC	False Colour Composite
OCP	Opencast Project
UGP	Underground Project
OB	Over Burden
GCP	Ground Control points
GIS	Geographic Information System
WGS-84	World Geodetic System
UTM	Universal Transverse Mercator

## GLOSSARY

Sl.	Term	Definition
1.	Land Reclamation	To manage, reclaim and restore mined out/ degraded land as close as possible to its original stage.
2.	Over Burden	The material that lies above the coal seam/ deposit
3.	Monitoring	A process of evaluation to check or keep record for a period of time.
4.	Opencast Coal Mine	Open-pit mining, also known as opencast mining, is a surface mining technique that extracts minerals from an open pit in the ground.
5.	Social Forestry	Social forestry is the management and protection of forests and afforestation of barren and deforested lands with the purpose of helping environmental, social and rural development. Plantation (Social/ Avenue or other) carried out outside mining area.
6.	Biological Reclamation	Plantation on Backfilled areas (Stabilised Internal Dumps)
7.	Technical Reclamation	Area under backfilling (Over burden dumped inside the mine voids) in mining area.
8.	Green Cover Generated	Total Plantation carried out in the lease area of Project. This includes Plantation on Backfill, Plantation on OB and Social Forestry.
9.	Leasehold Area	The area, for which lease is granted for the purpose of undertaking mining and allied operations.
10.	Excavated area	Mined out area which includes active mining, area under backfilling and plantation on backfilled areas
11.	Active Mining	Mining areas which include Coal Quarry, Advance Quarry, Quarry Filled with Water etc.
12.	Environmental Protection	It is the practice of protecting the natural environment by individuals, organizations and governments. Its objectives are to conserve natural resources and the existing natural environment and, where possible, to mitigate damage and reverse trends.
13.	Remedial Measure	Any measure or action required or undertaken to investigate, monitor, clean up, remove, treat, prevent, contain or otherwise remediate the presence or release of any hazardous substance or activity.
14.	Systematic Error	Every measurement differing from the true measurement in the same direction, and even by the same amount in some cases.



15.	Geometric Distortion	It refers to the improper positioning of any image with respect to their true geographic position when viewed in a properly scaled common image display plane.
16.	Land Use/ Cover Class	Land cover is what covers the surface of the earth and land use describes how the land is used.
17.	Accuracy	The closeness of agreement between a measured quantity value and a true quantity value.
18.	Environmental Clearance	Environmental Clearance (EC) for any developmental projects like coal mining projects etc. has been made mandatory by the Ministry of Environment, Forests and Climate Change (MoEF & CC) through its Notification issued on 27.01.1994 under the provisions of Environment (Protection) Act, 1986.
19.	Rectification and Geo-referencing	Geo-referencing is the assigning of absolute location of a data point or data points. Geo-rectification refers to the removal of geometric distortions between sets of data points, most often the removal of terrain, platform, and sensor induced distortions from remote sensing imagery.
20.	Image Enhancement	It is the process of modifying digital images so that the results are more suitable for processing or further image analysis.
21.	Training set selection	It is a portion of a data set used to fit or train a model for prediction or classification of values that are known in the training set, but unknown in other (future) data.
22.	Image Classification	It refers to the task of extracting information classes from a multiband raster image. The resulting raster from image classification can be used to create thematic maps.
23.	Temporal Changes	The 'temporal change' means the change in any entity with a period of time.
24.	Ground Truthing	Collection of primary/ basic information from ground realities for satellite image interpretation and thematic mapping.
25.	Cluster	Group of opencast and/ or underground mines clubbed together for administrative purposes.
26.	Arc GIS	GIS Software used for Map preparation
27.	ERDAS IMAGINE	Satellite Image Data Classification Software



## Central Mine Planning & Design Institute Ltd.

(A Subsidiary of Coal India Ltd.)

Gondwana Place, Kanke Road, Ranchi 834031, Jharkhand

Phone : (+91) 651 2230001, 2230002, 2230483, FAX (+91) 651 2231447, 2231851

Website : [www.cmpdi.co.in](http://www.cmpdi.co.in), Email : [cmpdihq@cmpdi.co.in](mailto:cmpdihq@cmpdi.co.in)